



Opening New Frontiers in Singlet Fission Research



Remco W. A. Havenith

Stratingh Institute for Chemistry,
Zernike Institute for Advanced Materials,
University of Groningen,
The Netherlands

Ghent Quantum Chemistry Group, Ghent University, Belgium



university of
groningen

faculty of science
and engineering

stratingh institute
for chemistry

Who contributed



university of
groningen



Ria Broer



Luis Enrique



Kathir




Coen de Graaf



university of
groningen



UNIVERSITAT
ROVIRA I VIRGILI



Tjerk Straatsma



university of
groningen

faculty of science
and engineering

stratingh institute
for chemistry

FOM Focus Group Groningen 'Next Generation Organic Photovoltaics'

- Aim:
 - Deliver the science for highly efficient, long-lived, and low-cost organic photovoltaic devices
- Challenge:
 - Charge separation at the donor/acceptor interface
- Approach:
 - Multi-disciplinary:
 - Material development
 - Physical characterisation (OPV device physics)
 - *Theoretical modelling*



university of
 groningen

faculty of science
 and engineering


stratingh institute
 for chemistry

Theoretical challenges

- Predict molecular properties that determine the dielectric properties of the interface
 - Dipole moments
 - Polarisability
- Modelling of the donor/acceptor interface
 - Molecular Dynamics simulations
 - Time scales of molecular motion
- Calculation of the excited states
 - Theoretical methods
 - Influence of molecular structure
 - Influence of the embedding using multiscale modelling
- Approximation of the electron/energy transfer rates




Singlet fission



SF: spin allowed radiationless process

M. B. Smith, J. Michl, *Chem. Rev.* **110** (2010), 6891

It is attractive to build the wavefunctions of the solid from state-specific molecular wavefunctions



university of
groningen

faculty of science
and engineering

stratingh institute
for chemistry

Singlet fission rate

- Fermi Golden rule in diabatic representation (Marcus theory)

$$k_{SF} \propto \left| \langle \Psi_f | H | \Psi_i \rangle \right|^2 = \left| \langle S_0 S_1 | H | {}^1 T T \rangle \right|^2$$

- Electronic coupling between diabatic states
 - Directly accessible with our non orthogonal CI approach

- Adiabatic representation: Non-adiabatic couplings (Landau-Zener model)


$$\langle \Psi_f | \frac{\partial}{\partial Q} | \Psi_i \rangle$$

- Potential energy surfaces and conical intersections/crossings

P.F. Barbara, T.J. Meyer, M.A. Ratner, *J. Phys. Chem.* **100** (1996), 13148
F. Bernardi, M. Olivucci, and M.A. Robb, *Chem. Soc. Rev.* **25** (1996), 321



Cluster approximation for solids



- Describe solid in terms of molecular wavefunctions
- Compute wavefunctions of each molecule for specific states (CASSCF)
- Form many-electron basis functions (S_0S_0 , S_0S_1 , 1TT , CT), each describing a particular combination of molecular states




Non orthogonal configuration interaction

- Describe wavefunctions of a cluster of molecules in terms of (localised) molecular many-electron basis functions (MEBF)
 - MEBFs are spin-adapted antisymmetrised products of molecular wavefunctions:

$$\Phi_{AB}^{KL} = A(\Phi_A^K \times \Phi_B^L)$$

- Molecular wavefunction can be any multiconfigurational wavefunction



Ψ^0 : Ground state

Ψ^1 : Singlet excited state

Ψ^T : Triplet excited state

$$\Phi_{AB}^{00} = A(\Psi_A^0 \times \Psi_B^0)$$

$$\Phi_{AB}^{10} = A(\Psi_A^1 \times \Psi_B^0)$$

$$\Phi_{AB}^{01} = A(\Psi_A^0 \times \Psi_B^1)$$

$$\Phi_{AB}^{TT} = A(\Psi_A^T \times \Psi_B^T)$$

\vdots

$$\Psi = c_1 \Phi_{AB}^{00} + c_2 \Phi_{AB}^{10} + c_3 \Phi_{AB}^{01} + c_4 \Phi_{AB}^{TT} + \dots$$




Non Orthogonal Configuration Interaction

- Wavefunction expanded as: $\Psi = \sum_{i=1}^N C_i \Phi_i$ with Φ_i a many-electron basis function ((MEBF) Slater determinant, or combination thereof)
- The orbitals ϕ_j in a MEBF are not orthogonal, making the many-electron MEBFs also not orthogonal: $\langle \Phi_i | \Phi_j \rangle = S_{ij}$
- The non orthogonality of the orbitals within one MEBF and of the orbitals in a different MEBF complicates the calculation of the required Hamiltonian matrix elements $\langle \Phi_i | H | \Phi_j \rangle$
- Solve $(\mathbf{H} - E\mathbf{S})(\mathbf{C}) = 0$ to get energies and Ψ (C_i 's)



Advantages of this NOCI

- Inclusion of orbital relaxation effects
- Inclusion of (static) correlation effects
- Short wavefunction expansions
- Chemical interpretability
 - Description of system in terms of predefined states



- **Con:** no simple Slater rules for the computation of matrix elements of the Hamilton operator in the MEBF basis



Computational Aspects of our NOCI approach

- Typical number of $|\Phi_{AB}^{KL}\rangle \sim 20$
 - H/S matrices contain ~ 210 elements of the type
 $\langle \Phi_{AB}^{KL} | H | \Phi_{AB}^{K'L'} \rangle$ and $\langle \Phi_{AB}^{KL} | \Phi_{AB}^{K'L'} \rangle$
- If $|\Phi_A^K\rangle$ contains ~ 500 determinants, then $|\Phi_A^K\rangle \times |\Phi_B^L\rangle = |\Phi_{AB}^{KL}\rangle \sim 2.5 \cdot 10^5$
- $\langle \Phi_{AB}^{KL} | H | \Phi_{AB}^{K'L'} \rangle = \sum_i \sum_j c_i c_j \langle \Delta_i | H | \Delta_j \rangle$
- Approximately 10^7 elements $\langle \Delta_i | H | \Delta_j \rangle$ have to be calculated for one matrix element $\langle \Phi_{AB}^{KL} | H | \Phi_{AB}^{K'L'} \rangle$
- Aim for high level of parallelism
- Easy to parallelize




Technical Aspects

- Evaluation of $\langle \Delta_i | H | \Delta_j \rangle$ with non orthogonal orbitals
 - $H_{ij} = \sum_{i,j} h_{ij} S^{(i,j)} + \sum_{i < k} \sum_{j < l} [(ij | kl) - (ik | jl)] S^{(i,j,k,l)}$
 - First and second order co-factors needed
- With corresponding orbitals, then $\langle c_i | d_j \rangle = \lambda_i \delta_{ij}$
and $S^{(i,i)} = \prod_{m \neq i} \lambda_m$ ($S^{(i,j)} = 0$ for $i \neq j$)
- No 4-index, but transform co-factors to common basis in which the corresponding orbitals c_i and d_i are expressed
- SVD and matrix multiplications
- Use GPUs



The GronOR code



- In collaboration with OLCF, based on the GNOME code
 - OpenACC for GPU off-loading
 - Master-worker model with task based load balancing
 - MPI parallelization with point-to-point non-blocking communication
 - Avoid global synchronization and global reduction operations
 - Fault resilient implementation




university of
 groningen

faculty of science
 and engineering

stratingh institute
 for chemistry

GronOR Master-Worker Process Layout



Each process group has the same number of worker processes

Each process group should have sufficient aggregate memory to hold all integrals:

One-electron integrals are duplicated

Two-electron integrals are distributed

Consequences of Node Faults: All processes on a failing node fail
If a worker process fails, the entire group to which it belongs will fail





university of
groningen

faculty of science
and engineering

stratingh institute
for chemistry

Case study I - Naphthalene dimer

Anti-symmetrised products of CAS(4,4) wavefunctions

				E (eV)
	Ψ_A^0	\times	Ψ_B^0	0.00
	Ψ_A^1	\times	Ψ_B^0	6.54
	Ψ_A^0	\times	Ψ_B^1	6.54
	Ψ_A^T	\times	Ψ_B^T	8.64


- Neglect of dynamical correlation (S_1 too high)
- Endoergic



Performance of GronOR

Benchmark run on Summit, requiring 112,867,800 matrix element evaluations

- Each node with 6 MPI ranks
- 1 GPU per rank
- Good scalability with number of nodes
- Performance improvement from GPU is **6.8x**



university of
groningen

faculty of science
and engineering

stratingh institute
for chemistry

Reduction of required 2-e⁻ integrals

- The MOs in the CASSCF wavefunction are expressed in N AOs
 - # 2-e⁻ integrals $\sim N^4/8$


- The n inactive + active MOs of all states of a molecule form a basis as well:

$$\{ \phi_1^{S_0}, \dots, \phi_n^{S_0}, \phi_1^{S_1}, \dots, \phi_n^{S_1} \}$$

- Redundant basis, eliminate linear dependencies, based on threshold ϵ , and transform MOs and 2-e⁻ integrals to new basis
- Dimension of new basis $m \ll N$



Case study II - Pyridine



570	148	136	110	98	72	54	48	#bf
46000	841	600	257	162	47	15	9.1	Integral size (MB)
2408	253	244	226	219/ 73	210	205	204	Time (s)




university of
groningen

faculty of science
and engineering

stratingh institute
for chemistry

CI threshold – remove contributions from configurations with small CI coefficients



1024		1024		64	1024*			Nodes
6238		3002		406	87			Time (s)



university of
groningen



faculty of science
and engineering

stratingh institute
for chemistry



*6 ranks/node

Singlet fission in tetracene

- Geometry dependence of coupling in solid
 - MD using DFTB (NVT)
 - Estimate coupling using DFT and NOCI



Effective coupling, DFT and NOCI




- Coupling heavily dependent on geometry
- AB/AC orientation best




Tetracene coupling

- Large electronic coupling in the AB orientation
- Molecules A and B get closer
 - Larger HOMO/HOMO and LUMO/LUMO overlap
- Strong mixing of CT states (30%) in local excited state on A and in ^1TT state
- Direct coupling: 5 meV
- CT mediated coupling: 32 meV



Correlation NOCI vs DFT



- Only weak correlation



university of
groningen

$\log(V_{\text{NOCI}})$

faculty of science
and engineering



stratingh institute
for chemistry

Conclusions and outlook

- Parallel NOCI program GronOR is working and ready to be used for interesting applications
 - Further optimizations are in progress
 - Better handling of CI lists
 - Inclusion of dynamical correlation and embedding effects
- Tetracene
 - CT states enhance the coupling
 - Coupling and CT state mixing heavily dependent on geometry
 - Static picture not always sufficient for predicting singlet fission
 - Weak correlation NOCI vs DFT couplings
- Further studies:
 - Cibalackrot and other SF molecules
 - Effects of functional groups on electronic coupling



Theoretical Chemistry Group



university of
groningen

faculty of science
and engineering

stratingh institute
for chemistry

Financial support

- OLCF for computer time on SUMMIT (ESP)
- FOM Focus Group 'Next Generation Organic Photovoltaics' (DIFFER)
- FOM/Shell
- ITN-EJD-TCCM (Horizon2020)



university of
 groningen

faculty of science
 and engineering

stratingh institute
 for chemistry



university of
 groningen

faculty of science
and engineering

stratingh institute
for chemistry